

ALEXANDROFF TYPE MANIFOLDS AND HOMOLOGY MANIFOLDS

V. TODOROV AND V. VALOV

ABSTRACT. One of the open questions in generalized Cantor manifolds is whether every Alexandroff manifold with respect to the class D_G^{n-2} of all spaces whose cohomological dimension $\dim_G$ is $\leq n - 2$, where G is an abelian group, is a Mazurkiewicz D_G^{n-2} -manifold. In our search for the answer of this question we introduce and investigate different types of connectedness between disjoint subsets of compacta, and prove that every strong K_G^n -manifold is a Mazurkiewicz D_G^{n-2} -manifold. We also establish a result implying that if X is arcwise connected complete metric space which is either a homology n -manifold over a group G or a product of n metric spaces, then X is a Mazurkiewicz arc n -manifold.

1. INTRODUCTION

Cantor manifolds were introduced by Urysohn [19] as a generalization of Euclidean manifolds. Recall that a space X is a *Cantor n -manifold* if any partition of X is of dimension $\geq n - 1$ [19] (a partition of X is a closed set $P \subset X$ such that $X \setminus P$ is the union of two open disjoint sets). In other words, X cannot be the union of two proper closed sets whose intersection is of covering dimension $\leq n - 2$. *Strong Cantor manifolds* is another specification of Cantor manifolds which was considered by Hadžiivanov [6]. Hadžiivanov and Todorov [8] introduced the class of *Mazurkiewicz n -manifolds*, which is a proper sub-class of the strong Cantor n -manifolds: X is a Mazurkiewicz n -manifold if for any disjoint closed massive subsets A and B of X (a massive subset of X is a set with non-empty interior in X), and any normally placed set M of X with $\dim M \leq n - 2$, there exists a continuum in $X \setminus M$ intersecting A and B (equivalently, no such M is cutting X between A and B). Here, M is a normally placed in X provided every two disjoint closed in M sets have disjoint open in X neighborhoods (for example, every F_σ -

2010 *Mathematics Subject Classification*. Primary 55M10, 55M15; Secondary 54F45, 54C55.

Key words and phrases. Cantor n -manifold, cohomological dimension, cohomology groups, homology n -manifold, V^n -continuum.

The second author was partially supported by NSERC Grant 261914-08.

subset of a normal space is normally placed in that space). The notion of a Mazurkiewicz n -manifold has its roots in the classical Mazurkiewicz theorem [13] that no region X in the Euclidean n -space can be cut by a subset M with $\dim M \leq n - 2$ in following sense: any two points from $X \setminus M$ can be join by a continuum $K \subset X \setminus M$.

But the strongest specification of Cantor manifolds is the notion of V^n -continua introduced by Alexandroff [1]: a compactum X is a V^n -continuum if for every two closed disjoint massive subsets X_0, X_1 of X there exists an open cover ω of X such that there is no partition P in X between X_0 and X_1 admitting an ω -map into a space Y with $\dim Y \leq n - 2$ ($f: P \rightarrow Y$ is said to be an ω -map if there exists an open cover γ of Y such that $f^{-1}(\gamma)$ refines ω).

The above notions are related as follows: strong Cantor n -manifolds are Cantor n -manifolds, every V^n -continuum is a Mazurkiewicz n -manifold and Mazurkiewicz n -manifolds are strong Cantor n -manifolds [8]. None of the above inclusions is reversible [9].

More general concepts of the above notions were considered in [9] and [16]. In particular, we are going to use the following two, where $\mathcal{C}$ is a class of topological spaces.

Definition 1.1. A space X is an *Alexandroff manifold with respect to $\mathcal{C}$* (br., *Alexandroff $\mathcal{C}$ -manifold*) if for every two closed, disjoint, massive subsets X_0, X_1 of X there exists an open cover ω of X such that no partition P in X between X_0 and X_1 admits an ω -map onto a space $Y \in \mathcal{C}$.

Definition 1.2. A space X is said to be a *Mazurkiewicz manifold with respect to $\mathcal{C}$* (br., *Mazurkiewicz $\mathcal{C}$ -manifold*) provided for every two closed, disjoint, massive subsets $X_0, X_1 \subset X$, and every set $F = \bigcup_{i=0}^{\infty} F_i \subset X$ with each $F_i \in \mathcal{C}$ being proper closed subset of X , there exists a continuum K in $X \setminus F$ joining X_0 and X_1 . If, in the above definition, X_0 and X_1 can be joined by an arc in $X \setminus F$, X is called a *Mazurkiewicz arc $\mathcal{C}$ -manifold*.

It is still unknown if every Alexandroff D_G^{n-2} -manifold is a Mazurkiewicz D_G^{n-2} -manifold, where D_G^{n-2} is the class of all spaces whose cohomological dimension $\dim_G$ is $\leq n - 2$. Searching for the answer of this question, we consider in Section 2 different types of connectedness between disjoint subsets of compacta. The starting point for our considerations in Section 3 was the following result of Krupski [10, Proposition 1.7]: *Let X be a locally compact locally connected separable metric space such that for all $k < n$ and $x \in X$ the singular homology groups $H_k(X, X \setminus x)$ are trivial. Then every open connected*

subset of $U \subset X$ is a Cantor n -manifold. We establish that, under the hypotheses of Krupski's result, every connected open subset of X is a Mazurkiewicz arc manifold with respect to the class of all spaces whose covering dimension is $\leq n - 2$. The same conclusion also holds for arcwise connected open subsets of a complete metric space which is a product of n metric spaces.

Finally, let us raise the following question (see Section 2 for the corresponding definitions):

Question 1.3. *Let X be an Alexandroff manifold with respect to the class D_G^{n-2} . Is it true that X is a Mazurkiewicz D_G^{n-2} -manifold? What about when (X, F) is a K_G^n -manifold or X is a V_G^n -continuum?*

2. ALEXANDROFF TYPES CONNECTEDNESS OF SPACES

In this section all spaces are assumed to be at least paracompact and G denotes an abelian group. For any space X the cohomological dimension $\dim_G X$ is the smallest integer n such that $H^{n+1}(X, A : G) = 0$ for all closed subsets $A \subset X$. Here, $H^{n+1}(X, A : G)$ denotes the Čech cohomology group of the pair (X, A) with coefficients from G . If ω is an open cover of X and $Z \subset X$ is closed, we denote by $|\omega|$ and $|\omega_Z|$ the nerves of ω and the system $\omega_Z = \{U \cap Z : U \in \omega\}$. For any such ω and $Z \subset X$ let $p_{\omega_Z}^* : H^k(|\omega|, |\omega_Z| : G) \rightarrow H^k(X, Z : G)$, $k \geq 0$, be the projection between the k -th cohomology groups, where $p_{\omega_Z} : (X, Z) \rightarrow (|\omega|, |\omega_Z|)$ is a map generated by a partition of unity subordinated to ω (we call such a map p_{ω_Z} to be natural). Further, $i_A : (A, B) \rightarrow (X, Z)$ denotes the embedding of the pair (A, B) into the pair (X, Z) . Everywhere in this section we suppose that the coefficients of the cohomology groups are from a fixed group G and omit the symbol for the group of coefficients.

Definition 2.1. Let P and Q be disjoint open subsets of a compactum X and $F \subset X$ closed. We say that the pair (X, F) is K_G^n -connected between P and Q if there exist an open cover ω of $Y = X \setminus (P \cup Q)$ such that the following condition holds for every partition C of X between P and Q : any natural map $p_{\omega_C} : (C, C \cap F) \rightarrow (|\omega_C|, |\omega_{C \cap F}|)$ generates a non-trivial homomorphism

$$p_{\omega_C}^* : H^{n-1}(|\omega_C|, |\omega_{C \cap F}|) \rightarrow H^{n-1}(C, C \cap F).$$

If, in the above situation, there exists also $e \in H^{n-1}(|\omega|, |\omega_{F \cap Y}|)$ such that $p_{\omega_C}^*(i_{\omega_C}^*(e)) \neq 0$ for every partition C in X between P and Q , the pair (X, F) is called *strongly K_G^n -connected between P and Q* . Further, (X, F) is said to be a K_G^n -manifold (resp., *strong K_G^n -manifold*) if it is

K_G^n -connected (resp., strongly K_G^n -connected) between any two open disjoint sets $P, Q \subset X$.

Definition 2.1 is justified by the following result which was actually established by Kuzminov [12]:

Theorem 2.2. *Every compactum X with $\dim_G X \geq n$ contains a pair (Y, F) of closed sets such that (Y, F) is a strong K_G^n -manifold.*

Similarly, the definition of Alexandroff $\mathcal{C}$ -manifold yields the following one:

Definition 2.3. Let P and Q be open subsets of a compactum X with $\overline{P} \cap \overline{Q} = \emptyset$. We say that X is $\mathcal{C}$ -connected in the sense of Alexandroff between P and Q (shortly, $A(\mathcal{C})$ -connected between P and Q) if there exists an open cover ω of $X \setminus (P \cup Q)$ such that no partition of X between P and Q admits an ω -map onto a space from the class $\mathcal{C}$.

Proposition 2.4. *Let P and Q be open subsets of a compactum X with $\overline{P} \cap \overline{Q} = \emptyset$ and $F \subset X$ closed. If the pair (X, F) is K_G^n -connected between P and Q , then X is $A(D_G^{n-2})$ -connected between P and Q , where D_G^{n-2} is the class of spaces of dimension $\dim_G \leq n - 2$.*

Proof. According to Definition 2.1, we can find an open cover ω of $Y = X \setminus (P \cup Q)$ satisfying the following condition: for every partition C in X between P and Q and a natural map $p_{\omega_C}: (C, C \cap F) \rightarrow (|\omega_C|, |\omega_{C \cap F}|)$ there exists an element $e_C \in H^{n-1}(|\omega_C|, |\omega_{C \cap F}|)$ with $p_{\omega_C}^*(e_C) \neq 0$.

Suppose there exists a partition C of X between P and Q admitting an ω -map $g: C \rightarrow T$ onto a compactum T with $\dim_G T \leq n - 2$. Thus, we can find a finite open cover τ of T such that $\nu = g^{-1}(\tau)$ is refining ω . Let $p_\nu: (C, C \cap F) \rightarrow (|\nu|, |\nu_{C \cap F}|)$ and $p_\tau: (T, g(C \cap F)) \rightarrow (|\tau|, |\tau_{g(C \cap F)}|)$ be natural maps. Obviously, the function $V \in \tau \rightarrow g^{-1}(V) \in \nu$ provides a simplicial homeomorphism $g_\nu^\tau: (|\tau|, |\tau_{g(C \cap F)}|) \rightarrow (|\nu|, |\nu_{C \cap F}|)$. Then the maps p_ν and $g_\tau = g_\nu^\tau \circ p_\tau \circ g$ are homotopic. Hence,

$$(1) \quad p_\nu^* = g^* \circ p_\tau^* \circ (g_\nu^\tau)^*.$$

Because $H^{n-1}(T, g(C \cap F)) = 0$ (recall that $\dim_G T \leq n - 2$), the homomorphism $g^*: H^{n-1}(T, g(C \cap F)) \rightarrow H^{n-1}(C, C \cap F)$ is trivial. Then, by (1), so is the homomorphism $p_\nu^*: H^{n-1}(|\nu|, |\nu_{C \cap F}|) \rightarrow H^{n-1}(C, C \cap F)$. On the other hand, since ν refines ω , we can find a map $\varphi_\nu: (|\nu|, |\nu_{C \cap F}|) \rightarrow (|\omega_C|, |\omega_{C \cap F}|)$ such that p_{ω_C} and $\varphi_\nu \circ p_\nu$ are homotopic. Therefore, $p_{\omega_C}^* = p_\nu^* \circ \varphi_\nu^*$. Since $p_{\omega_C}^*(e_C) \neq 0$ for some $e_C \in H^{n-1}(|\omega_C|, |\omega_{C \cap F}|)$, $p_\nu^*(\varphi_\nu^*(e_C)) \neq 0$, which contradicts the triviality of p_ν^* .

Therefore, X is $A(D_G^{n-2})$ -connected between P and Q . □

Corollary 2.5. *A compactum X is an Alexandroff D_G^{n-2} -manifold provided (X, F) is a K_G^n -manifold for some closed set $F \subset X$.*

Definition 2.6. Let X, P, Q and F be as in Definition 2.1. We say that the pair (X, F) is V_G^n -connected between P and Q if there exists an open cover ω of $X \setminus (P \cup Q)$ such that $g^*: H^{n-1}(T, g(C \cap F)) \rightarrow H^{n-1}(C, C \cap F)$ is non-trivial for any partition C of X between P and Q and any surjective ω -map $g: C \rightarrow T$. The pair (X, F) is called a *relative V_G^n -continuum* provided (X, F) is V_G^n -connected between any two open sets $P, Q \subset X$ with disjoint closures.

When F is the empty set, the above definition provides V_G^n -continua introduced by Stefanov [15]. The proof of Proposition 2.3 yields the following stronger conclusion:

Corollary 2.7. *If a pair (X, F) is K_G^n -connected between two open sets $P, Q \subset X$ with disjoint closures, then (X, F) is V_G^n -connected between P and Q . In particular, every K_G^n -manifold (X, F) is a relative V_G^n -continuum.*

There is an interesting analogy between V_G^n -continua and relative V_G^n -continua. It follows from [12] and Corollary 2.7 that every compactum X with $\dim_G X = n$ contains a relative V_G^n -continuum. On the other hand, any compactum X with $H^n(X; G) \neq 0$ contains a V_G^n -continuum (see [15] for finite-dimensional metric X , and [20] for any compact X).

Another type of connectedness is inspired by Mazurkiewicz $\mathcal{C}$ -manifolds, where $\mathcal{C}$ is a given class of spaces.

Definition 2.8. Let $P, Q \subset X$ be open subsets of X with disjoint closures. The space X is said to be $M(\mathcal{C})$ -connected between P and Q if for every set $F \subset X$ with $F \in \mathcal{C}$ there exists a continuum $K \subset X \setminus F$ connecting P and Q .

It is well known [8] that any compact space X which is $A(D^{n-2})$ -connected between two open sets $P, Q \subset X$ with disjoint closures is M -connected between P and Q with respect to the class of all normally placed $(n-2)$ -dimensional subsets of X (here D^{n-2} is the class of all spaces of covering dimension $\dim \leq n-2$). We don't know if this holds when the covering dimension is replaced by the cohomological dimension $\dim_G$. But this is true if instead of $A(D^{n-2})$ -connectedness the stronger K_G^n -connectedness of X is assumed.

Theorem 2.9. *Let the pair (X, F) be strongly K_G^n -connected between two open disjoint sets $P, Q \subset X$. Then for every Lindelöf normally placed subset $M \subset X$ with $\dim_G M \leq n-2$ there exists a continuum $K \subset X \setminus M$ connecting P and Q .*

Proof. Suppose there exists a Lindelöf normally placed subset $M \subset X$ with $\dim_G M \leq n-2$ such that any continuum K connecting P and Q meets M . According to [9, Lemma 2.5], there are open sets $P_1, Q_1 \subset X$ such that $\overline{P_1} \subset P$, $\overline{Q_1} \subset Q$ and if K is a continuum in X joining $\overline{P_1}$ and $\overline{Q_1}$, then $K \cap M \setminus (P_1 \cup Q_1) \neq \emptyset$. We are going to show this is not true, which will complete the proof.

Let ω be a finite open cover of $Y = X \setminus (P_1 \cup Q_1)$ and e an element of $H^{n-1}(|\omega|, |\omega_{F \cap Y}|)$, both satisfying the requirements from Definition 2.1 with P and Q replaced by P_1 and Q_1 , respectively. Obviously, $M_1 = M \setminus (P_1 \cup Q_1)$ is closed in M . So, M_1 is also Lindelöf and normally placed in Y . Moreover, $\dim_G M_1 \leq n-2$, which implies $H^{n-1}(M_1, M_1 \cap F) = 0$. Consequently, there exists an open cover γ of M_1 refining ω and a simplicial map

$$\pi_\omega^\gamma : (|\gamma|, |\gamma_{M_1 \cap F}|) \rightarrow (|\omega_{M_1}|, |\omega_{M_1 \cap F}|) \text{ such that } (\pi_\omega^\gamma)^*(i_{M_1}^*(e)) = 0,$$

where i_{M_1} is the embedding $(|\omega_{M_1}|, |\omega_{M_1 \cap F}|) \hookrightarrow (|\omega|, |\omega_F|)$. Since M_1 is Lindelöf, we may assume that $\gamma = \{U_i : i = 1, 2, \dots\}$ is countable. Next, let $\gamma_1 = \{A_i : i = 1, 2, \dots\}$ be a system of closed in M_1 sets covering $M_1 \cap F$ such that $A_i \subset U_i \cap F$ for all i . Using the proof of [4, Theorem 3.1.1], we find an open in M_1 system $\{L_i : i = 1, 2, \dots\}$ such that $\lambda_1 = \{\overline{L_i}^{M_1} : i = 1, 2, \dots\}$ is a swelling of γ_1 with $\overline{L_i}^{M_1} \subset U_i$, $i \geq 1$, where $\overline{L_i}^{M_1}$ is the closure of L_i in M_1 . Recall that λ_1 is a swelling of γ_1 if each $\overline{L_i}^{M_1}$ contains A_i and for any finite set of indices $i_1, \dots, i_s$ we have $\bigcap_{k=1}^{k=s} \overline{L_{i_k}}^{M_1} \neq \emptyset$ iff $\bigcap_{k=1}^{k=s} A_{i_k} \neq \emptyset$. Consider the Lindelöf space $M_1 \setminus L$, where $L = \bigcup_{i \geq 1} L_i$. If $M_1 \setminus L$ is non-empty, there exists a countable system $\lambda_2 = \{B_i : i = 1, 2, \dots\}$ of closed subsets of M_1 such that λ_2 covers $M_1 \setminus L$, λ_2 refines γ and all B_i are contained in $M_1 \setminus F$. In case $M_1 \setminus L = \emptyset$, λ_2 is the empty family. Obviously, $\lambda = \lambda_1 \cup \lambda_2$ is a closed countable cover of M_1 refining γ . Let $\lambda = \{\Lambda_i : i = 1, 2, \dots\}$. Using again the proof of [4, Theorem 3.1.1], we find an open in M_1 swelling $\theta = \{W_i : i = 1, 2, \dots\}$ of λ such that θ refines γ . Moreover, we can insist that $W_i \cap F = \emptyset$ provided $\Lambda_i \cap F = \emptyset$.

Claim. *There exists an open in Y swellings $\tilde{\sigma} = \{V_i : i = 1, 2, \dots\}$ of λ refining ω such that $\Lambda_i \subset \overline{V_i} \cap M_1 \subset W_i$ for all i .*

We follow the proof of Lemma 4.1.2 from [17]. Obviously, it suffices to construct by induction open in Y swellings $\sigma_m = \{V_1, \dots, V_m\}$ of $\{\Lambda_1, \dots, \Lambda_m\}$, $m \geq 1$, such that

- $\sigma_{m+1} = \sigma_m \cup \{V_{m+1}\}$;
- $\Lambda_k \subset V_k \cap M_1 \subset \overline{V_k} \cap M_1 \subset W_k$, $k \geq 1$;
- each V_i is contained in an element of ω .

Suppose we already constructed σ_m for some m . Since Λ_{m+1} and $M \setminus W_{m+1}$ are closed disjoint subsets of M_1 and M_1 is normally placed in Y , there are disjoint open in Y sets V'_{m+1} and $\tilde{V}_{m+1}$ with $\Lambda_{m+1} \subset V'_{m+1}$ and $M_1 \setminus W_{m+1} \subset \tilde{V}_{m+1}$. Let E be the union of all intersections $\overline{V}_{i_1} \cap \dots \cap \overline{V}_{i_s}$, $1 \leq i_k \leq m$, such that $\bigcap_{k=1}^{k=s} \Lambda_{i_k} \cap \Lambda_{m+1} = \emptyset$. We claim that $E \cap \Lambda_{m+1} = \emptyset$. Indeed, otherwise $\bigcap_{k=1}^{k=p} \overline{V}_{j_k} \cap \Lambda_{m+1} \neq \emptyset$ for some indices $j_1, \dots, j_p$ such that $j_k \leq m$ and $\bigcap_{k=1}^{k=p} \Lambda_{j_k} \cap \Lambda_{m+1} = \emptyset$. Then

$$\bigcap_{k=1}^{k=p} \overline{V}_{j_k} \cap \Lambda_{m+1} \subset \bigcap_{k=1}^{k=p} \overline{V}_{j_k} \cap M \cap \Lambda_{m+1} \subset \bigcap_{k=1}^{k=p} W_{j_k} \cap W_{m+1} \neq \emptyset.$$

Since $\{W_i : i = 1, 2, \dots\}$ is a swelling of λ , this implies that $\bigcap_{k=1}^{k=p} \Lambda_{j_k} \cap \Lambda_{m+1} \neq \emptyset$, a contradiction. To finish the inductive step, observe that λ refines ω . So, there exists $O \in \omega$ containing Λ_{m+1} . We set $V_{n+1} = V'_{m+1} \cap O \setminus E$, and $\sigma_{m+1} = \sigma_m \cup \{V_{m+1}\}$. It is easily seen that σ_{m+1} is a swelling of $\{\Lambda_i : i = 1, 2, \dots, m+1\}$, which completes the proof of the claim.

Recall that θ is a refining of γ . Moreover, any refining function between θ and γ provides a simplicial map

$$\pi_\gamma^\theta : (|\theta|, |\theta_{M_1 \cap F}|) \rightarrow (|\gamma_{M_1}|, |\gamma_{M_1 \cap F}|) \text{ such that } (\pi_\omega^\theta)^*(i_{M_1}^*(e)) = 0,$$

where π_ω^θ is the composition $\pi_\omega^\gamma \circ \pi_\gamma^\theta$. Let $\theta_j = \{W_i : \Lambda_i \in \lambda_j\}$, $j = 1, 2$. Obviously, $\theta = \theta_1 \cup \theta_2$. Consider the open in Y system

$$\sigma = \{V_i \setminus F : W_i \in \theta_2\} \cup \{V_i : W_i \in \theta_1\} \text{ and let } W = \bigcup_{\Sigma \in \sigma} \Sigma.$$

It follows from our construction that $\sigma = \{\Sigma_i : i = 1, 2, \dots\}$ has the following property: for any indices $i_1, \dots, i_s$ we have

$$\bigcap_{k=1}^{k=s} \Sigma_{i_k} \neq \emptyset \text{ iff } \bigcap_{k=1}^{k=s} W_{i_k} \neq \emptyset \text{ and } \bigcap_{k=1}^{k=s} \Sigma_{i_k} \cap F \neq \emptyset \text{ iff } \bigcap_{k=1}^{k=s} W_{i_k} \cap F \neq \emptyset.$$

Therefore, there exists a simplicial homeomorphism ξ between the nerves $(|\theta|, |\theta_{M_1 \cap F}|)$ and $(|\sigma|, |\sigma_{W \cap F}|)$. So, ξ^{-1} is also a simplicial homeomorphism from $(|\sigma|, |\sigma_{W \cap F}|)$ onto $(|\theta|, |\theta_{M_1 \cap F}|)$. Then for the simplicial map π below we have

$$\pi = i_{M_1} \circ \pi_\omega^\theta \circ \xi^{-1} : (|\sigma|, |\sigma_{W \cap F}|) \rightarrow (|\omega|, |\omega_{F \cap Y}|), \quad \pi^*(e) = 0.$$

Suppose the set W contains a partition C of X between $\overline{P}_1$ and $\overline{Q}_1$ and consider the following diagram, where i and i_C are natural inclusions:

$$\begin{array}{ccccc}
 (|\sigma_C|, |\sigma_{F \cap C}|) & \xrightarrow{i} & (|\sigma|, |\sigma_{W \cap F}|) & \xrightarrow{\pi} & (|\omega|, |\omega_{F \cap Y}|) \\
 \uparrow p_{\sigma_C} & & & & \uparrow p_{\omega} \\
 (C, C \cap F) & \xrightarrow{i_C} & (Y, F \cap Y) & &
 \end{array}$$

It follows from our construction that for every $x \in C$ (resp., $x \in C \cap F$) both points $\pi(i(p_{\sigma_C}(x)))$ and $p_{\omega}(i_C(x))$ belong to a simplex of the nerve $|\omega|$ (resp. $|\omega_{F \cap Y}|$). So, the maps $\pi \circ i \circ p_{\sigma_C}$ and $p_{\omega} \circ i_C$ are homotopic and we have the commutative diagram:

$$\begin{array}{ccccc}
 H^{n-1}(|\omega|, |\omega_{F \cap Y}|) & \xrightarrow{\pi^*} & H^{n-1}(|\sigma|, |\sigma_{W \cap F}|) & \xrightarrow{i^*} & H^{n-1}(|\sigma_C|, |\sigma_{C \cap F}|) \\
 \downarrow p_{\omega}^* & & & & \downarrow p_{\sigma_C}^* \\
 H^{n-1}(Y, F \cap Y) & \xrightarrow{i_C^*} & H^{n-1}(C, C \cap F) & &
 \end{array}$$

Because $\pi^*(e) = 0$, the last diagram implies

$$(2) \quad i_C^*(p_{\omega}^*(e)) = 0.$$

On the other hand, the following diagram

$$\begin{array}{ccc}
 H^{n-1}(|\omega|, |\omega_{F \cap Y}|) & \xrightarrow{i_{\omega_C}^*} & H^{n-1}(|\omega_C|, |\omega_{C \cap F}|) \\
 \downarrow p_{\omega}^* & & \downarrow p_{\omega_C}^* \\
 H^{n-1}(Y, F \cap Y) & \xrightarrow{i_C^*} & H^{n-1}(C, C \cap F)
 \end{array}$$

yields

$$(3) \quad i_C^*(p_{\omega}^*(e)) = p_{\omega_C}^*(i_{\omega_C}^*(e)) = 0.$$

Combining (2) and (3), we obtain $p_{\omega_C}^*(i_{\omega_C}^*(e)) = 0$, a contradiction.

Hence, W does not contain any partition of X between $\overline{P}_1$ and $\overline{Q}_1$. Choose an open set $\tilde{W}$ such that both sets $\overline{P}_1 \setminus \tilde{W}$ and $\overline{Q}_1 \setminus \tilde{W}$ are non-empty and $\tilde{W} \cap Y = W$. Suppose that $X \setminus \tilde{W}$ is not connected between

$\overline{P}_1 \setminus \tilde{W}$ and $\overline{Q}_1 \setminus \tilde{W}$, i.e. $X \setminus \tilde{W} = A \cup B$ with A, B being disjoint closed sets such that $\overline{P}_1 \setminus \tilde{W} \subset A$ and $\overline{Q}_1 \setminus \tilde{W} \subset B$. Then the sets $B' = B \cup \overline{Q}_1$ and $A' = A \cup \overline{P}_1$ are disjoint and closed in X whose union is a proper subset of X . So, there is a partition C in X between A' and B' . Obviously, C is a partition between $\overline{P}_1$ and $\overline{Q}_1$ which is contained in W , a contradiction. Hence, $X \setminus \tilde{W}$ is connected between $\overline{P}_1 \setminus \tilde{W}$ and $\overline{Q}_1 \setminus \tilde{W}$. This implies (see [11, §47, Theorem 3]) the existence of a continuum $K \subset X \setminus \tilde{W}$ connecting $\overline{P}_1$ and $\overline{Q}_1$. Finally, since $M_1 \subset \tilde{W}$, $K \subset X \setminus M_1$. This contradicts the fact that any continuum connecting $\overline{P}_1$ and $\overline{Q}_1$ should meet M_1 . $\square$

Corollary 2.10. *Every strong K_G^n -manifold is a Mazurkiewicz D_G^{n-2} -manifold.*

3. HOMOLOGY MANIFOLDS AND MAZURKIEWICZ MANIFOLDS

In this section we are going to show that some homological properties of a metric space X imply that X is a *Mazurkiewicz arc n -manifold* in the following sense: If M is an F_σ -subset of X with $\dim M \leq n - 2$, then any two massive disjoint sets A and B can be joined by an arc.

Everywhere in this section we consider singular homology groups reduced in dimension zero with coefficients in a given group G (if G is not written then the coefficients are integers). The following notion introduced by Toruńczyk [18] is well known: A closed subset A of a space X is said to be a Z_n -set in X if for any map $f: \mathbb{I}^n \rightarrow X$ and any open cover ω of X there is a map $g: \mathbb{I}^n \rightarrow X$ which is ω -close to f (i.e., for all $x \in \mathbb{I}^n$ both points $f(x)$ and $g(x)$ belong to some element of ω). A homological counterpart of this notion was defined by Banach-Cauty-Karashev [3]: A closed subset $A \subset X$ is called a G -homological Z_n -set in X if $H_k(U, U \setminus A; G) = 0$ for all $k \leq n$ and all open $U \subset X$. It follows from the excision axiom that a point $x \in X$ is a G -homological Z_n -set in X provided $H_k(X, X \setminus x; G) = 0$ for all $k \leq n$.

Further, let us remind the definition of the separating dimension $t(X)$ of a space X introduced by Steinke [14] and its transfinite extension $\text{trt}(X)$ given by Arenas-Chatyrko-Puertas [2]: $\text{trt}(X) = -1$ iff $X = \emptyset$; $\text{trt}(X) \leq \alpha$ for an ordinal α if any closed set $B \subset X$ with containing at least two points can be separated by a closed set $P \subset B$ with $\text{trt}(P) < \alpha$. When $\text{trt}(X)$ is an integer, then $\text{trt}(X) = t(X)$. Moreover, $t(X) \leq \text{ind}(X)$ [14]. Hence, for metrizable X we have $t(X) \leq \text{ind}(X) \leq \dim X$.

Theorem 3.1. *Let X be a complete metric space and M be an F_σ -subset of X such that $\text{trt}(M) \leq n - 2$ and $H_k(X, X \setminus x; G) = 0$ for all*

$k \leq n - 1$ and all $x \in M$. Suppose $P, Q \subset X$ are open sets which can be joined by an arc in X . Then there is an arc in $X \setminus M$ joining P and Q . In particular, any arcwise connected open subset of X is a Mazurkiewicz arc n -manifold provided that $H_k(X, X \setminus x; G) = 0$ for all $k \leq n - 1$ and $x \in X$.

Proof. Let M be a countable union of closed sets M_i , $i = 1, 2, \dots$. Since each $x \in M$ is a G -homological Z_{n-1} -point in X and $\text{trt}(M_i) \leq n-2$, we can apply Theorem 4.3 from [3] stating that if A is a closed subset of a space X such that $\text{trt}(A) = m$ and all $a \in A$ are G -homological Z_{n+m} -points in X , then A is a G -homological Z_{n+m} -set in X . Therefore, any one of the sets M_i is a G -homological Z_1 -set in X . Then, by [3, Theorem 3.2(6)], M_i are Z_1 -sets in X . Consequently, the spaces $C_i = C(\mathbb{I}, X \setminus M_i)$ are dense (and obviously, open) in the space $C(\mathbb{I}, X)$ of all continuous maps from $\mathbb{I}$ into X equipped with the compact-open topology. Finally, since $C(\mathbb{I}, X)$ is complete, $\bigcap C_i$ is also dense in $C(\mathbb{I}, X)$. Because P and Q can be joined by an arc in X , there exists a map $f: \mathbb{I} \rightarrow X$ with $f(0) \in P$ and $f(1) \in Q$. Therefore, the map f can be approximated by maps $f': \mathbb{I} \rightarrow X \setminus M$. Since P and Q are open, there exists $g: \mathbb{I} \rightarrow X \setminus M$ with $g(0) \in P$ and $g(1) \in Q$.

The second half follows from the first one and the following observations: if $U \subset X$ is open, then the excision axiom implies $H_k(U, U \setminus x; G) = 0$ for all $k \leq n - 1$ and $x \in U$; moreover, $\text{trt}(M) \leq \dim M$ for any set $M \subset X$. $\square$

Below, by an n -manifold over G we mean a metric space X such that for every $x \in X$ we have $H_k(X, X \setminus x; G) = 0$ if $k \neq n$ and $H_n(X, X \setminus x; G) = G$.

Corollary 3.2. *Let X be an arcwise connected complete metric space. In each of the following cases any arcwise connected open subset of X is a Mazurkiewicz arc n -manifold:*

- (1) X is a homology n -manifold over a group G ;
- (2) X is a product of n metric spaces X_i , $1 \leq i \leq n$.

Proof. The first item follows directly from Theorem 3.1. To prove the second one, consider the exact sequence for every i and $x \in X_i$

$$H_0(X_i \setminus x; G) \rightarrow H_0(X_i; G) \rightarrow H_0(X_i, X_i \setminus x; G) \rightarrow 0.$$

Since X_i is arcwise connected (as an image of X), $H_0(X_i; G) = 0$. So, $H_0(X_i, X_i \setminus x; G) = 0$. Then, by [3, Theorem 3.2 and Theorem 6.1(2)], $H_k(X, X \setminus x; G) = 0$ for every $x \in X$ and $k \leq n - 1$. Finally, Theorem 3.1 completes the proof. $\square$

Next theorem shows that in some situations the space $X \setminus M$ from Theorem 3.1 is arcwise connected.

Theorem 3.3. *Let X be a connected and locally connected complete metric space and M be an F_σ -set in X with $\dim M \leq n - 2$. Then $X \setminus M$ is arcwise connected in each of the following cases:*

- (1) X is a homology n -manifold over a group G ;
- (2) X is a product of n metric spaces.

Proof. Since X is connected and locally connected, it is arcwise connected and locally arcwise connected. Let $M = \bigcup M_i$ with each M_i being closed in X . It follows from the proofs of Theorem 3.1 and Corollary 3.2 that in any of the two cases M_i are Z_1 -sets in X . For any two points $a, b \in X \setminus M$ consider the set-valued map $\Phi: \mathbb{I} \rightarrow X$ defined by $\Phi(0) = a$, $\Phi(1) = b$ and $\Phi(t) = X$ for each $t \in (0, 1)$. It is easily seen that Φ is lower semi-continuous and for every $x \in X$ and its neighborhood U there is a neighborhood $V \subset U$ of x with the following property: if $t \in [0, 1]$ and x_1, x_2 are two points from $\Phi(t) \cap V$ there is an arc in $\Phi(t) \cap U$ joining x_1 and x_2 . Then, by [5, Theorem 1.2], Φ admits a continuous selection $g: \mathbb{I} \rightarrow X \setminus M$. Obviously, $g(\mathbb{I})$ is an arc in $X \setminus M$ joining a and b . $\square$

Finally, we provide some conditions for Cantor manifolds to be Mazurkiewicz manifolds.

Proposition 3.4. *Suppose X is a connected complete metric space possessing a base $\mathcal{B}$ of open sets such that $\overline{U}$ is a Cantor manifold with respect to a given class $\mathcal{C}$ for every $U \in \mathcal{B}$. Then $X \setminus M$ is arcwise connected for every F_σ -subset $M = \bigcup M_i$ of X with $M_i \in \mathcal{C}$, $i \geq 1$.*

Proof. According to a result of Hadžiivanov-Hamamdžiev [7, Theorem 1], $X \setminus M$ is connected. Moreover, by the same result, all $U \setminus M$, $U \in \mathcal{B}$, are connected. Hence, $X \setminus M$ is connected and locally connected. Because $X \setminus M$ is complete (as a G_δ -subset of X), it is arcwise connected. $\square$

REFERENCES

- [1] P. S. Alexandroff, *Die Kontinua (V^p) - eine Verschärfung der Cantorschen Mannigfaltigkeiten*, Monatshefte für Math. **61** (1957), 67–76 (German).
- [2] F. Arenas, V. Chatyrko and M. Puertas, *Transfinite extension of Steinke's dimension*, Acta Math. Hung. **88** (2000), 105–112.
- [3] T. Banach, R. Cauty and A. Karashev, *On homotopical and homological Z_n -sets*, Topology Proc. **38** (2011), 29–82.
- [4] R. Engelking, *Theory of dimensions: Finite and Infinite*, Heldermann Verlag, Lemgo, 1995.

- [5] V. Gutev and V. Valov, *Dense families of selections and finite-dimensional spaces*, Set-Valued Analysis **11** (2003), 373–391.
- [6] N. Hadžiivanov, *Strong Cantor manifolds*, C. R. Acad. Bulgare Sci. **30** (1977), 1247–1249 (Russian).
- [7] N. Hadžiivanov and A. Hamamdžiev, *A method of P.S. Urysohn for the proof that an F_σ -set does not cut*, Serdica **3** (1977), 130–135 (in Russian).
- [8] N. Hadžiivanov and V. Todorov, *On non-Euclidean manifolds*, C. R. Acad. Bulgare Sci. **33** (1980), 449–452 (Russian).
- [9] A. Karashev, P. Krupski, V. Todorov and V. Valov, *Generalized Cantor manifolds and homogeneity*, Houston J. Math. **38** (2012), no. 2, 583–609.
- [10] P. Krupski, *On the disjoint $(0, n)$ -cells property for homogeneous ANR's*, Colloq. Math. **66** (1993), no. 1, 77–84.
- [11] K. Kuratowski, *Topology*, vol. II, Academic Press, New York; PWN-Polish Scientific Publishers, Warsaw, 1968.
- [12] V. Kuz'minov, *On V^n continua*, Dokl. Akad. Nauk SSSR **139** (1961), 24–27 (Russian).
- [13] S. Mazurkiewicz, *Sur les ensembles de dimension faible*, Fund. Math. **13** (1929), 210–217.
- [14] G. Steinke, *A new dimension by means of separating sets*, Arch. Math. **40** (1983), 273–282.
- [15] S. Stefanov, *A cohomological analogue of V^n -continua and a theorem of Mazurkiewicz*, Serdica **12** (1986), no. 1, 88–94 (in Russian).
- [16] V. Todorov and V. Valov, *Generalized Cantor manifolds and indecomposable continua*, Questions and Answers in Gen. Topolol. **30** (2012), 93–102.
- [17] V. Todorov, , PhD Thesis, Sofia University, 1979.
- [18] H. Toruńczyk, *Concerning locally homotopy negligible sets and characterization of l_2 -manifolds*, Fund. Math. **101:2** (1978), 93–110.
- [19] P. Urysohn, *Memoire sur les multiplicites cantoriennes*, Fund. Math. **7** (1925), 30–137.
- [20] V. Valov, *Homogeneous ANR spaces and Alexandroff manifolds*, preprint.

DEPARTMENT OF MATHEMATICS, UACG, 1 H. SMIRNENSKI BLVD., 1046 SOFIA, BULGARIA

E-mail address: vtt-fte@uacg.bg

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, NIPISSING UNIVERSITY, 100 COLLEGE DRIVE, P.O. BOX 5002, NORTH BAY, ON, P1B 8L7, CANADA

E-mail address: veskov@nipissingu.ca